Table 5.12b Adjusted Patient Survival, <u>Deceased Donor</u> ECD Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3	Months		1 Year		5	5 Years		-	10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.
Total												
All	3,291	97.0%	0.4%	3,291	92.8%	0.8%	8,790	71.0%	1.1%	14,123	40.9%	0.8%
Age at Tx												
1-5 Years	0	-	-	o	-	-	0	-	-	1	+	+
6-11 Years	0	-	-	0	-	-	2	+	+	9	90.0%	9.5%
12-17 Years	1	+	+	1	+	+	4	+	+	27	91.9%	5.6%
18-34 Years	62	98.2%	1.8%	62	98.2%	1.8%	275	86.8%	2.4%	684	73.1%	2.4%
35-49 Years	340	98.5%	0.7%	340	96.6%	1.0%	1,269	83.6%	1.4%	2,585	61.7%	1.4%
50-64 Years	1,682	97.6%	0.4%	1,682	93.3%	0.6%	4,329	72.7%	1.0%	6,917	41.9%	1.2%
65+ Years	1,206	95.5%	0.6%	1,206	90.1%	0.9%	2,911	61.3%	1.4%	3,900	23.3%	1.9%
Recipient Ethnicity/Race												
White	1,576	97.0%	0.5%	1,576	91.8%	1.0%	4,223	68.5%	1.4%	7,031	38.6%	1.1%
African American	1,071	96.4%	0.7%	1,071	92.9%	1.0%	2,800	71.3%	1.5%	4,377	38.4%	1.6%
Hispanic/Latino	363	98.3%	0.7%	363	94.9%	1.2%	1,008	77.3%	2.0%	1,567	47.6%	2.5%
Asian	238	98.1%	0.9%	238	95.4%	1.4%	620	74.9%	2.7%	938	55.1%	3.3%
Other/Multi-race	43	96.5%	2.5%	43	94.7%	3.0%	139	76.9%	5.3%	210	41.9%	6.8%
Recipient Gender												
Female	1,207	97.2%	0.6%	1,207	93.6%	1.0%	3,232	72.8%	1.4%	5,331	44.7%	1.3%
Male	2,084	96.9%	0.5%	2,084	92.4%	1.0%	5,558	69.9%	1.3%	8,792	38.6%	1.1%

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

⁽⁺⁾ = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category. Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12b (Continued) Adjusted Patient Survival, <u>Deceased Donor</u> ECD Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3	3 Months			1 Year		Ę	5 Years			10 Years		
	(Tx 2006 - 2007)			(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)			
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Total All	3,291	97.0%	0.4%	3,291	92.8%	0.8%	8,790	71.0%	1.1%	14,123	40.9%	0.8%	
Primary Diagnosis													
Glomerular Diseases	467	98.6%	0.6%	467	94.9%	1.2%	1,379	79.1%	1.7%	2,501	53.0%	1.9%	
Diabetes Hypertensive	1,094	96.3%	0.7%	1,094	91.0%	1.4%	2,857	63.0%	1.7%	4,302	27.6%	1.49	
Nephrosclerosis	975	96.5%	0.7%	975	93.0%	1.2%	2,352	71.7%	1.6%	3,435	42.2%	1.89	
Polycystic Kidneys Tubular and Interstitial	274	98.7%	0.6%	274	95.6%	1.3%	748	82.5%	1.9%	1,291	58.8%	2.59	
Diseases Renovascular & Other	131	98.0%	1.2%	131	95.2%	1.9%	360	68.7%	3.6%	628	46.4%	3.59	
Vascular Diseases Congenital, Rare Familial, & Metabolic	108	96.2%	1.9%	108	89.6%	3.2%	382	68.6%	3.3%	711	35.2%	3.19	
Disorders	19	100.0%	0.0%	19	88.1%	7.7%	61	84.1%	5.7%	112	62.6%	10.19	
Neoplasms	10	+	+	10	+	+	37	57.4%	11.7%	54	30.0%	11.19	
Other	211	96.5%	1.3%	211	92.6%	1.9%	609	71.9%	2.7%	1,082	43.9%	2.69	
Unknown	2	+	+	2	+	+	5	80.9%	16.3%	7	85.0%	13.39	

Source: OPTN/SRTR Data as of May 4, 2009.

Notes:

^{(+) =} Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12b (Continued) Adjusted Patient Survival, <u>Deceased Donor</u> ECD Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3 Months				1 Year		5	Years		10 Years (Tx 1997 - 2007)			
	(Tx 2	2006 - 20	007)	(Tx 2	2006 - 20	007)	(Tx 2002 - 2007)						
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Total													
All	3,291	97.0%	0.4%	3,291	92.8%	0.8%	8,790	71.0%	1.1%	14,123	40.9%	0.8%	
Transplant Center State													
Alabama	25	92.4%	5.1%	25	88.6%	6.2%	104	70.5%	5.7%	225	32.2%	5.39	
Arkansas	24	96.1%	3.8%	24	92.1%	5.3%	50	54.1%	13.4%	67	25.0%	10.39	
Arizona	46	98.0%	1.9%	46	94.1%	3.3%	107	80.1%	5.4%	151	65.2%	6.59	
California	339	98.1%	0.7%	339	94.1%	1.4%	1,010	72.1%	2.1%	1,662	44.9%	2.39	
Colorado	22	100.0%	0.0%	22	96.4%	3.5%	65	82.5%	5.2%	94	29.9%	10.19	
Connecticut	28	97.0%	3.0%	28	93.9%	4.2%	66	72.7%	6.3%	94	59.2%	6.59	
District of Columbia	46	96.0%	2.8%	46	94.0%	3.4%	88	81.1%	5.0%	141	23.6%	6.89	
Delaware	2	+	+	2	+	+	2	+	+	2	+	ĺ -	
Florida	182	95.9%	1.5%	182	91.8%	2.1%	478	65.6%	3.8%	774	34.9%	3.59	
Georgia	52	93.0%	3.4%	52	87.8%	4.5%	169	77.2%	3.8%	288	43.2%	6.19	
Hawaii	27	100.0%	0.0%	27	96.3%	3.5%	55	70.7%	9.3%	96	59.0%	7.39	
Iowa	15	100.0%	0.0%	15	94.4%	5.3%	48	74.0%	10.3%	84	38.9%	8.79	
Illinois	132	96.3%	1.7%	132	88.8%	2.9%	399	69.5%	3.2%	617	41.5%	3.89	
Indiana	12	100.0%	0.0%	12	94.0%	5.6%	71	73.9%	8.0%	96	54.1%	9.69	
Kansas	12	+	+	12	+	+	22	48.0%	20.4%	26	60.7%	16.9	
Kentucky	34	+	+	34	+	+	79	67.2%	7.8%	137	43.4%	8.29	
Louisiana	52	97.7%	2.3%	52	95.4%	3.2%	129	82.2%	4.7%	184	44.7%	7.79	
Massachusetts	101	97.6%	1.4%	101	96.8%	1.6%	254	77.2%	3.6%	365	39.5%	5.79	
Maryland	116	98.2%	1.2%	116	96.5%	1.8%	331	61.2%	4.1%	655	33.1%	3.3	
Maine	2	+	+	2	+	+	10	+	+	33	63.9%	9.49	
Michigan	104	95.9%	1.9%	104	90.9%	2.8%	229	66.6%	4.6%	343	39.5%	5.39	
Minnesota	32	+	+	32	+	+	85	85.4%	6.0%	192	42.2%	5.69	
Missouri	54	96.7%	2.3%	54	91.8%	3.6%	104	87.5%	3.7%	182	46.7%	5.89	
Mississippi	2	+	+	2	+	+	3	58.8%	27.9%	13	23.3%	12.69	
North Carolina	111	95.6%	2.0%	111	92.0%	2.7%	281	74.1%	4.2%	423	45.7%	5.19	
North Dakota	7	100.0%	0.0%	7	87.1%	10.8%	36	52.1%	9.9%	58	29.5%	11.59	
Nebraska	24	100.0%	0.0%	24	92.1%	5.3%	48	85.0%	5.9%	74	32.9%	12.79	
New Hampshire	14	100.0%	0.0%	14	93.7%	6.0%	22	83.1%	7.6%	29	55.0%	15.09	
New Jersey	101	94.3%	2.3%	101	86.6%	3.6%	333	66.1%	3.8%	528	40.3%	4.99	
New Mexico	13	100.0%	0.0%	13	93.7%	5.9%	35	91.6%	4.6%	62	47.1%	12.79	
Nevada	4	+	+	4	+	+	14	72.3%	13.4%	32	29.0%	11.09	

(Continued)

Source: OPTN/SRTR Data as of May 4, 2009.

Notes

^{(+) =} Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.

Table 5.12b (Continued) Adjusted Patient Survival, <u>Deceased Donor</u> ECD Kidney Transplants Survival at 3 Months, 1 Year, 5 Years, and 10 Years

	3	3 Months			1 Year		5 Years			10 Years			
	(Tx 2006 - 2007)			(Tx 2	(Tx 2006 - 2007)			(Tx 2002 - 2007)			(Tx 1997 - 2007)		
	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	N	%	Std. Err.	
Transplant Center State													
New York	393	96.7%	1.0%	393	93.4%	1.4%	899	69.5%	2.7%	1,356	43.7%	2.5%	
Ohio	123	97.1%	1.5%	123	91.8%	2.5%	330	67.9%	3.7%	569	37.2%	3.6%	
Oklahoma	30	94.2%	3.9%	30	91.3%	4.8%	71	71.3%	6.6%	117	50.4%	8.1%	
Oregon	38	93.8%	3.5%	38	91.7%	4.0%	124	72.5%	4.9%	192	59.1%	6.0%	
Pennsylvania	322	97.6%	0.8%	322	92.3%	1.6%	881	70.2%	2.3%	1,391	37.9%	2.7%	
Puerto Rico	23	100.0%	0.0%	23	95.6%	4.2%	49	89.0%	5.4%	64	27.0%	16.1%	
Rhode Island	4	+	+	4	+	+	34	73.5%	8.1%	61	36.4%	13.8%	
South Carolina	27	96.8%	3.1%	27	96.8%	3.1%	63	57.4%	10.4%	103	39.1%	7.4%	
South Dakota	12	100.0%	0.0%	12	87.3%	8.1%	63	67.9%	8.0%	75	17.9%	10.8%	
Tennessee	69	92.8%	3.2%	69	88.5%	4.0%	172	69.2%	5.3%	286	32.2%	5.7%	
Texas	228	96.4%	1.3%	228	93.6%	1.8%	599	76.3%	2.7%	905	45.6%	3.4%	
Utah	13	94.1%	5.4%	13	88.2%	7.7%	24	71.3%	11.3%	44	45.2%	11.0%	
Virginia	113	99.1%	0.9%	113	91.6%	2.8%	263	66.8%	4.7%	368	34.4%	6.1%	
Vermont	8	91.1%	8.1%	8	91.1%	8.1%	21	79.8%	11.4%	28	76.6%	8.9%	
Washington	38	100.0%	0.0%	38	97.4%	2.5%	126	74.1%	5.4%	259	48.5%	4.9%	
Wisconsin	111	98.4%	1.1%	111	92.2%	2.5%	326	67.5%	3.7%	522	36.7%	4.4%	
West Virginia	4	+	+	4	+	+	18	86.8%	8.4%	56	46.2%	9.2%	

Source: OPTN/SRTR Data as of May 4, 2009.

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on kidney and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Survival rates are adjusted to the characteristics of the 3 month and 1 year cohort. See Technical Notes for details.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether kidneys were allocated under ECD system. Non-ECDs include all other donors, including SCDs and DCDs.